

**Experimenteller Nachweis der Existenz von**

# **F**ar**b**l**a**d**u**ng

**Dariusz Miśkowiec, 9.03.2009**

# Farbladung – Ladung der starken Wechselwirkung



**Gravitation**



**Schwache Wechselwirkung**



**Elektromagnetische WW**



**Starke Wechselwirkung**

# Hinweis 1: Existenz von $\Delta^{++}$ und $\Omega^{-}$

(Gell-Mann 1972, Fritzsche 1973)

$$\Omega^{-} \sim \underbrace{\mathbf{s}^{\uparrow} \mathbf{s}^{\uparrow} \mathbf{s}^{\uparrow}}_{S=3/2} \underbrace{\psi(\mathbf{x}_1, \mathbf{x}_2, \mathbf{x}_3)}_{L=0 \text{ symmetrisch}}$$

Pauli Prinzip  $\rightarrow$  die 3 Quarks dürfen nicht identisch sein

Farbladung bietet einen Ausweg

$$\Omega^{-} = \frac{1}{\sqrt{6}} \varepsilon_{\alpha\beta\gamma} \mathbf{s}_{\alpha}^{\uparrow} \mathbf{s}_{\beta}^{\uparrow} \mathbf{s}_{\gamma}^{\uparrow} \psi(\mathbf{x}_1, \mathbf{x}_2, \mathbf{x}_3)$$

# Hadronen invariant unter $SU_c(3)$



- ☉ Farbladungszustände (z.B.): **rot**, **grün**, **blau**
- ☉  $SU_c(3)$ : Rotationen im Farbraum
- ☉ Hadronen: Farb-Singulett-Zustände
- ☉ Baryonen: **qqq** + **qqq** + **qqq** - **qqq** - **qqq** - **qqq**
- ☉ Mesonen: **q $\bar{q}$**  + **q $\bar{q}$**  + **q $\bar{q}$**

"Farbe" - weil weiß = **rot** + **grün** + **blau**

...aber nur für Menschen



# Hinweis 2: $e^+e^-$ Annihilation




Erwartung ohne Farbe:

$$R = \frac{\sigma(e^+e^- \rightarrow \text{Hadronen})}{\sigma(e^+e^- \rightarrow \text{Myonen})} = Q_u^2 + Q_d^2 + \dots = \left\{ \begin{array}{l} 4/9, \text{ u} \\ 5/9, \text{ ud} \\ 6/9, \text{ uds} \\ 10/9, \text{ udsc} \\ 11/9, \text{ udsb} \\ 15/9, \text{ alle Quarks} \end{array} \right.$$


Erwartung mit Farbe:

$$R = Q_u^2 + Q_u^2 + Q_u^2 + Q_d^2 + \dots \quad (3 \times \text{mehr})$$




**Messung: konsistent mit Farbe**

# Hinweis 3: Verzweigungsverhältnisse im $\tau$ -Zerfall



Erwartung	ohne Farbe	mit Farbe	PDG
$\tau^- \rightarrow e^-, \nu_e$	33%	20%	17.9%
$\tau^- \rightarrow \mu^-, \nu_\mu$	33%	20%	17.4%
$\tau^- \rightarrow ud$	33%	60%	


# Hinweis 4: Drell-Yan-Prozess



$$\frac{d^2\sigma}{dM^2 dx_F} = \frac{4\pi\alpha^2}{3M^4} \left(\frac{1}{3}\right) \sum_i \frac{x_1 x_2}{x_1 + x_2} Q_i^2 [f_{q_i}^{h_1}(x_1) f_{\bar{q}_i}^{h_2}(x_2) + f_{\bar{q}_i}^{h_1}(x_1) f_{q_i}^{h_2}(x_2)]$$

**Nur 1/3 von den Quark-Antiquark  
Paaren haben die passenden Farben**






**Rechnung mit Farbfaktor stimmt mit dem Experiment überein**

# Hinweis 5: $\pi^0$ - Zerfallsbreite

K. Huang, Quarks, Leptons and Gauge Fields  
Goity et al., Phys. Rev. D 66:076014, 2002




$$\Gamma(\omega) \sim 3 [ (2/3)^2 - (1/3)^2 ]$$

$$\Gamma(\pi^0 \rightarrow \gamma\gamma) = (\alpha/F_\pi)^2 (m_\pi/4/\pi)^3 + \dots$$

$$\Gamma(\pi^0 \rightarrow \gamma\gamma) = 8.10 \text{ eV} \pm 1.0\%$$

# Primakoff-Effekt

$$d\sigma/d\Omega \sim \Gamma(\pi^0 \rightarrow \gamma\gamma)$$




$$\tau(\pi^0) = (8.2 \pm 0.24) \times 10^{-17} \text{ s}$$

$$\Gamma(\pi^0 \rightarrow \gamma\gamma) = 8.1(2) \text{ eV}$$

# PrimEx-Experiment, JLab, 2008

Preliminary  $\pi^0$  Photoproduction yield ( $^{12}\text{C}$ , crystal only)



Tue Apr 8 09:38:35 2008

**Rechnung mit Farbfaktor stimmt mit dem Experiment überein**

# Farbladung nachgewiesen

1.  $\Delta^{++}$  und  $\Omega^{-}$  existieren
2.  $e^+e^- \rightarrow$  Hadronen 3 x höher
3.  $\tau \rightarrow$  Hadronen 3 x höher
4.  $q\bar{q}$ -Annihilation in DY 3 x niedriger
5.  $\pi^0 \rightarrow \gamma\gamma$  Rate 9 x höher

**BACKUP**

# Farbladung – Ladung der starken Wechselwirkung

## PROPERTIES OF THE INTERACTIONS

Property \ Interaction	Gravitational	Weak (Electroweak)	Electromagnetic	Strong	
				Fundamental	Residual
Acts on:	Mass – Energy	Flavor	Electric Charge	Color Charge	See Residual Strong Interaction Note
Particles experiencing:	All	Quarks, Leptons	Electrically charged	Quarks, Gluons	Hadrons
Particles mediating:	Graviton (not yet observed)	$W^+$ $W^-$ $Z^0$	$\gamma$	Gluons	Mesons
Strength relative to electromag for two u quarks at: for two protons in nucleus	$10^{-41}$	0.8	1	25	Not applicable to quarks
	$10^{-41}$	$10^{-4}$	1	60	
	$10^{-36}$	$10^{-7}$	1	Not applicable to hadrons	20

## BOSONS

force carriers  
spin = 0, 1, 2, ...

Unified Electroweak spin = 1		
Name	Mass GeV/c <sup>2</sup>	Electric charge
$\gamma$ photon	0	0
$W^-$	80.4	-1
$W^+$	80.4	+1
$Z^0$	91.187	0

Strong (color) spin = 1		
Name	Mass GeV/c <sup>2</sup>	Electric charge
$g$ gluon	0	0

# FERMIONS

matter constituents  
spin = 1/2, 3/2, 5/2, ...

Leptons spin = 1/2			Quarks spin = 1/2		
Flavor	Mass GeV/c <sup>2</sup>	Electric charge	Flavor	Approx. Mass GeV/c <sup>2</sup>	Electric charge
$\nu_e$ electron neutrino	$<1 \times 10^{-8}$	0	<b>u</b> up	0.003	2/3
<b>e</b> electron	0.000511	-1	<b>d</b> down	0.006	-1/3
$\nu_\mu$ muon neutrino	$<0.0002$	0	<b>c</b> charm	1.3	2/3
<b><math>\mu</math></b> muon	0.106	-1	<b>s</b> strange	0.1	-1/3
$\nu_\tau$ tau neutrino	$<0.02$	0	<b>t</b> top	175	2/3
<b><math>\tau</math></b> tau	1.7771	-1	<b>b</b> bottom	4.3	-1/3

## Mesons $q\bar{q}$

Mesons are bosonic hadrons.  
There are about 140 types of mesons.

Symbol	Name	Quark content	Electric charge	Mass GeV/c <sup>2</sup>	Spin
$\pi^+$	pion	$u\bar{d}$	+1	0.140	0
$K^-$	kaon	$s\bar{u}$	-1	0.494	0
$\rho^+$	rho	$u\bar{d}$	+1	0.770	1
$B^0$	B-zero	$d\bar{b}$	0	5.279	0
$\eta_c$	eta-c	$c\bar{c}$	0	2.980	0



## Baryons $qqq$ and Antibaryons $\bar{q}\bar{q}\bar{q}$

Baryons are fermionic hadrons.  
There are about 120 types of baryons.

Symbol	Name	Quark content	Electric charge	Mass $\text{GeV}/c^2$	Spin
<b>p</b>	proton	<b>uud</b>	1	0.938	1/2
<b><math>\bar{p}</math></b>	anti-proton	<b><math>\bar{u}\bar{u}\bar{d}</math></b>	-1	0.938	1/2
<b>n</b>	neutron	<b>udd</b>	0	0.940	1/2
<b><math>\Lambda</math></b>	lambda	<b>uds</b>	0	1.116	1/2
<b><math>\Omega^-</math></b>	omega	<b>sss</b>	-1	1.672	3/2

# Standard Model of FUNDAMENTAL PARTICLES AND INTERACTIONS


The Standard Model summarizes the current knowledge in Particle Physics. It is the quantum theory that includes the theory of strong interactions (quantum chromodynamics or QCD) and the unified theory of weak and electromagnetic interactions (electroweak). Gravity is included on this chart because it is one of the fundamental interactions even though not part of the "Standard Model."

## FERMIONS

matter constituents  
spin = 1/2, 3/2, 5/2, ...

Leptons spin = 1/2		
Flavor	Mass GeV/c <sup>2</sup>	Electric charge
$\nu_e$ electron neutrino	<1×10 <sup>-8</sup>	0
e electron	0.000511	-1
$\nu_\mu$ muon neutrino	<0.0002	0
$\mu$ muon	0.106	-1
$\nu_\tau$ tau neutrino	<0.02	0
$\tau$ tau	1.7771	-1

Quarks spin = 1/2		
Flavor	Approx. Mass GeV/c <sup>2</sup>	Electric charge
u up	0.003	2/3
d down	0.006	-1/3
c charm	1.3	2/3
s strange	0.1	-1/3
t top	175	2/3
b bottom	4.3	-1/3



If the protons and neutrons in this picture were 10 cm across, then the quarks and electrons would be less than 0.1 mm in size and the entire atom would be about 10 km across.

## BOSONS

force carriers  
spin = 0, 1, 2, ...

Unified Electroweak spin = 1		
Name	Mass GeV/c <sup>2</sup>	Electric charge
$\gamma$ photon	0	0
W <sup>-</sup>	80.4	-1
W <sup>+</sup>	80.4	+1
Z <sup>0</sup>	91.187	0

Strong (color) spin = 1		
Name	Mass GeV/c <sup>2</sup>	Electric charge
g gluon	0	0

**Color Charge**  
Each quark carries one of three types of "strong charge," also called "color charge." These charges have nothing to do with the colors of visible light. There are eight possible types of color charge for gluons. Just as electrically-charged particles interact by exchanging photons, in strong interactions color-charged particles interact by exchanging gluons. Leptons, photons, and W and Z bosons have no strong interactions and hence no color charge.

### Quarks Confined in Mesons and Baryons

One cannot isolate quarks and gluons; they are confined in color-neutral particles called **hadrons**. This confinement (binding) results from multiple exchanges of gluons among the color-charged constituents. As color-charged particles (quarks and gluons) move apart, the energy in the color-force field between them increases. This energy eventually is converted into additional quark-antiquark pairs (see figure below). The quarks and antiquarks then combine into hadrons; these are the particles seen to emerge. Two types of hadrons have been observed in nature: **mesons**  $q\bar{q}$  and **baryons**  $qqq$ .

### Residual Strong Interaction

The strong binding of color-neutral protons and neutrons to form nuclei is due to residual strong interactions between their color-charged constituents. It is similar to the residual electrical interaction that binds electrically neutral atoms to form molecules. It can also be viewed as the exchange of mesons between the hadrons.

## PROPERTIES OF THE INTERACTIONS

Baryons $qqq$ and Antibaryons $\bar{q}\bar{q}\bar{q}$					
Baryons are fermionic hadrons. There are about 120 types of baryons.					
Symbol	Name	Quark content	Electric charge	Mass GeV/c <sup>2</sup>	Spin
p	proton	uud	1	0.938	1/2
$\bar{p}$	anti-proton	$\bar{u}\bar{u}\bar{d}$	-1	0.938	1/2
n	neutron	udd	0	0.940	1/2
$\Lambda$	lambda	uds	0	1.116	1/2
$\Omega^-$	omega	sss	-1	1.672	3/2

Property	Interaction	Weak	Electromagnetic	Strong	
		(Electroweak)			Fundamental
Acts on:	Mass - Energy	Flavor	Electric Charge	Color Charge	See Residual Strong Interaction Note
Particles experiencing:	All	Quarks, Leptons	Electrically charged	Quarks, Gluons	Hadrons
Particles mediating:	Graviton (not yet observed)	W <sup>+</sup> W <sup>-</sup> Z <sup>0</sup>	$\gamma$	Gluons	Mesons
Strength relative to electromag for two u quarks at:	10 <sup>-18</sup> m	0.8	1	25	Not applicable to quarks
	3×10 <sup>-17</sup> m	10 <sup>-4</sup>	1	60	
for two protons in nucleus	10 <sup>-36</sup>	10 <sup>-7</sup>	1	Not applicable to hadrons	20




Mesons $q\bar{q}$					
Mesons are bosonic hadrons. There are about 140 types of mesons.					
Symbol	Name	Quark content	Electric charge	Mass GeV/c <sup>2</sup>	Spin
$\pi^+$	pion	$u\bar{d}$	+1	0.140	0
K <sup>-</sup>	kaon	$s\bar{u}$	-1	0.494	0
$\rho^+$	rho	$u\bar{d}$	+1	0.770	1
B <sup>0</sup>	B-zero	$d\bar{b}$	0	5.279	0
$\eta_c$	eta-c	$c\bar{c}$	0	2.980	0

### Matter and Antimatter

For every particle type there is a corresponding antiparticle type, denoted by a bar over the particle symbol (unless + or - charge is shown). Particle and antiparticle have identical mass and spin but opposite charges. Some electrically neutral bosons (e.g., Z<sup>0</sup>,  $\gamma$ , and  $\eta_c = c\bar{c}$ , but not K<sup>0</sup> = d $\bar{s}$ ) are their own antiparticles.

### Figures

These diagrams are an artist's conception of physical processes. They are **not** exact and have **no** meaningful scale. Green shaded areas represent the cloud of gluons or the gluon field, and red lines the quark paths.



### The Particle Adventure

Visit the award-winning web feature *The Particle Adventure* at <http://ParticleAdventure.org>

This chart has been made possible by the generous support of:

U.S. Department of Energy  
U.S. National Science Foundation  
Lawrence Berkeley National Laboratory  
Stanford Linear Accelerator Center  
American Physical Society, Division of Particles and Fields  
**BURLE INDUSTRIES, INC.**

©2000 Contemporary Physics Education Project. CPEP is a non-profit organization of teachers, physicists, and educators. Send mail to: CPEP, MS 50-308, Lawrence Berkeley National Laboratory, Berkeley, CA, 94720. For information on charts, text materials, hands-on classroom activities, and workshops, see:

<http://CPEPweb.org>

# Sehen

- 🌐 **Stäbchen:** hell/dunkel, empfindlich, Bewegung, Rand, blau
- 🌐 **Zapfen:** Farben, Auflösung, rot
  
- 🌐 **Mensch:** Trichromat – 3 Typen von Zapfen, rot, grün, blau
- 🌐 **Hund:** Dichromat – 2 Typen
- 🌐 **Vogel:** vielleicht 4 (UV)